

Influence of High Protein Digestibility Sorghums on Free Amino Nitrogen (FAN) Production during Malting and Mashing

Luke Mugode^{1,2}, Ostilio R. Portillo³, Dirk B. Hays³, Lloyd W. Rooney^{4,5}
and John R. N. Taylor^{1,*}

ABSTRACT

J. Inst. Brew. 117(3), 422–426, 2011

In sorghum brewing, obtaining sufficient Free Amino Nitrogen (FAN) for rapid and complete fermentation remains a problem due to the high proportions of unmalted sorghum used and the poor digestibility of wet-heat treated sorghum protein. Sorghum mutant lines with high protein digestibility have been developed through breeding. These high protein digestibility sorghums (HPDS) have protein bodies with villi-like borders that apparently facilitate protease access. This work investigated FAN production from HPDS when malted and mashed, to assess their potential for use in sorghum brewing to improve wort FAN levels. When malted, HPDS contained substantially higher levels of FAN than normal protein digestibility sorghums (NPDS), 32 mg/100 g malt more. However, when the HPDS were mashed either as malt, or as grain or malt plus exogenous proteases, FAN production during mashing was not substantially higher than with NPDS subjected to the same treatments, only 6, 6–18 and 9–13 mg/100 g grain or malt, respectively. This is probably due to wet-heat induced cross-linking of the kafirin proteins reducing their susceptibility to proteolysis. Notwithstanding this, HPDS could be very useful for improving FAN levels in sorghum brewing if they are malted.

Key words: exogenous proteases, free amino nitrogen (FAN), high protein digestibility sorghums (HPDS), malt, mashing, normal protein digestibility sorghums (NPDS), sorghum.

INTRODUCTION

In brewing, adequate levels of free amino acids and short peptides, collectively referred to as Free Amino Nitrogen (FAN) are required as a yeast nitrogen source for rapid and complete fermentation^{13,17}. When brewing with

sorghum this is a particular issue as the grists often comprise a low proportion of malt relative to unmalted adjunct or even 100% unmalted sorghum. Recent work on improving FAN production in sorghum brewing has focused on the use of exogenous proteases^{7,10,12}. However, it appears that to achieve sufficient proteolysis of the sorghum proteins, excessively high levels of exogenous proteases may be required. This can be attributed to the poor digestibility of the sorghum kafirin storage protein³, when it has been subjected to wet-heat treatment, as occurs in the brewing process.

Weaver and co-workers at Purdue University identified mutant sorghum lines with improved protein digestibility²³. These high protein digestibility sorghums (HPDS) were developed by crossing normal sorghum lines with a high-lysine sorghum mutant. More recently, research at Texas A&M University has focused on breeding to improve the agronomic and end-use quality of the HPDS, and determination of quantitative trait loci (QTLs) for the HPD trait²⁴.

This work therefore investigated FAN production from HPDS when malted and mashed, with the objective of assessing their potential for use in sorghum brewing to improve wort FAN levels.

MATERIALS AND METHODS

Materials

Sorghum cultivars. Normal protein digestibility sorghum (NPDS) parent (96GCPOB124) (Germinative Energy (GE) at 72 h 75%) (ICC), HPDS parent (P851171) (GE 92%) and HPDS progeny (04CS11278X 851171/96 GCP0124) (GE 90%) were produced at the experimental farm of Texas A&M University, College Station, Texas, USA. An unrelated NPDS cultivar (NK8828) (GE 89%) from the Agricultural Research Council, Potchefstroom, South Africa was included as control. All the sorghum lines were white, tan-plant types.

Proteases. Commercial proteolytic enzymes Flavourzyme 500 MG (an aminoprotease with 500 leucine aminopeptidase units/g) and Bioprotease P Conc. (an acid protease with 400,000 haemoglobin units/g) were kindly donated by Novozymes SA (Marlboro, South Africa) and Kerry Biosciences (Johannesburg, South Africa), respectively.

¹University of Pretoria, Department of Food Science, Private Bag X20, Hatfield 0028, South Africa.

²National Institute for Scientific and Industrial Research, P.O. Box 3101058, Lusaka 15302, Zambia.

³Texas A&M University, Cereal Genetics Laboratory, Department of Soil & Crop Sciences, College Station, TX 77843-2474, USA.

⁴Texas A&M University, Cereal Quality Laboratory, Department of Soil & Crop Sciences, College Station, TX 77843-2474, USA.

⁵Extraordinary Professor, University of Pretoria.

*Corresponding author. E-mail: John.taylor@up.ac.za

Table I. Grain characteristics and effects of malting on protein digestibilities of normal protein digestibility sorghum cultivars (NPDS) and on high protein digestibility sorghum (HPDS) cultivars^a.

Sorghum cultivar	Raw grain					Malt			
	Single kernel hardness	Thousand kernel weight	Endosperm texture ^b	Protein (% dry basis)	Lysine (g/100 g protein)	Protein digestibility (%)	Protein (% dry basis)	Protein digestibility (%)	Increase in protein digestibility with malting (%)
HPDS parent	33.5	21.6	5	10.1	3.9	65.8 ² ± 4.1	8.7	78.8 ³ ± 0.6	19.8
HPDS progeny	20.3	18.8	5	10.9	3.4	73.7 ² ± 3.1	9.0	82.9 ² ± 1.6	12.4
Means HPDS	26.9	20.2	5	10.5	3.7	69.8	8.9	80.9	16.1
NPDS parent	79.2	27.1	2	13.8	2.0	51.5 ¹ ± 1.5	12.1	67.5 ¹ ± 1.6	31.1
NPDS control	77.8	25.3	2	7.7	1.6	52.2 ¹ ± 2.3	6.9	71.8 ² ± 1.0	37.5
Means NPDS	78.5	26.2	2	10.4	1.8	51.9	9.5	69.7	34.3
Grand means	52.7	23.2	3.5	10.6	2.7	60.9	9.2	75.3	25.2

^a All results are means of at least two experiments. Means with different number superscripts within the columns differ significantly ($p < 0.05$).

^b Where 1 and 2 are corneous, 3 and 4 are intermediate, and 5 is floury.

Malting

The sorghum grains were malted according to agreed standard laboratory methods²⁰. Malt was dried at 50°C, after which the external roots and shoots were removed by rubbing in a coarse mesh bag so that roots and shoots were broken off and fell through the mesh. The sorghum grain and malt were milled using a laboratory-scale hammer mill type (Falling Number AB, Huddinge, Sweden) fitted with a 1.6 mm opening screen. The flour was stored in ziplock-type polyethylene bags at approximately 6°C until required.

Mashing

Ten grams of milled grain or malt was weighed into a 250 mL pre-weighed Erlenmeyer flask. Twenty mL tap water pre-heated to approximately 55°C and 1 mL calcium chloride solution (200 ppm) were added and the contents of the flask mixed thoroughly. The temperature of the mixture was kept at 55°C in a shaking water bath. In those treatments where the addition of exogenous proteases was investigated, 1 mL enzyme solution prepared to give a ratio of 1:100 (w/w) enzyme to flour was added and then mixed. The contents were mashed at 55°C for 45 min at the natural pH of the mash, approximately pH 5.8. At the end of mashing, supernatant was removed for FAN analysis.

Analyses

Grain physical characteristics. Three replicates of 300 sorghum kernels were each used to estimate grain hardness and 1,000 kernel weight using a Single Kernel Characterization System, SKCS 4100 (Perten Instruments, Springfield, IL). Grain endosperm texture was estimated by visual examination of longitudinal sectioned half kernels¹⁸.

Germinative energy (GE). Germinative Energy was determined according to ICC Draft Standard 174⁹.

Transmission electron microscopy (TEM). Sorghum grains were sectioned and then prepared for and viewed by TEM as described¹². Protein bodies in the sub-aleurone layers of the normal and high protein digestibility sorghums were compared.

Protein (N × 6.25). Protein was determined by a Dumas procedure¹.

Lysine. Lysine was determined by acid hydrolysis of the protein into its constituent amino acids, followed by

reverse-phase HPLC², using fluorescence detection. Human serum albumin was used as a control and norvaline and sarcosine were used as the internal standards.

In vitro protein digestibility. Protein digestibility was determined on the sorghum grain and malt samples using the pepsin method of Hamaker et al.⁸, where the solubilisation of the protein is measured after incubation at 37°C and pH 2.0 for 2 h. After incubation, the digest is centrifuged, the supernatant discarded, and the quantity of protein (the insoluble residue) is determined. Protein digestibility is calculated as protein solubilised (total protein in grain or malt minus residual protein) divided by total protein, and expressed as a percentage.

FAN. FAN was determined by the European Brewery Convention ninhydrin assay using glycine as the standard⁵ and expressed as mg FAN/100 g sorghum grain or malt.

Statistical analysis

All experiments were performed at least twice and closely agreeing replicate results were obtained. The data were analysed using one way analysis of variance and the means separated using Fischer's Least Significant Difference test at $p < 0.05$.

RESULTS AND DISCUSSION

The grain properties of the HPDS types were considerably different from those of the NPDS types. The NPDS were much harder, had much higher 1000 kernel weight (Table I), and had mainly a corneous type endosperm (Fig. 1). In contrast, not only were the HPDS softer and with a low 1000 kernel weight (Table I), the endosperms were also completely floury (Fig. 1). This confirms earlier reports the HPD trait in maize and sorghum tends to be associated with a soft endosperm²².

With regard to endosperm ultrastructure, TEM revealed that the surfaces of the protein bodies of the HPDS were irregular, with villi-like margins (indicated by arrows), compared to the protein bodies of the NPDS, which were essentially spherical with smooth surfaces (Fig. 2). It has been proposed that these invaginations in the surface of the protein bodies of the HPDS mutants apparently offer easier accessibility to proteolytic enzymes for digestion of the storage protein in the interior of the protein bodies¹⁵, which is mainly alpha-kafirin¹⁶.

With regard to protein related properties, while the protein content of the grains of two HPDS cultivars was in




Fig. 1. Endosperm texture of sorghum types. (A). NPDS parent, (B) NPDS, (C) HPDS parent, and (D) HPDS progeny.




Fig. 2. Transmission electron micrographs of protein bodies of different sorghum types. (a) NPDS (control), and (b) HPDS (progeny). PB – Protein Bodies, Arrows – Villi-like margins of protein bodies.

the same range as the NPDS, they had at least a 70% higher lysine content (Table I), resulting from their high-lysine mutant origin²³. As expected, the protein digestibilities of the HPDS were also higher than the NPDS, some 26–42% higher. This was higher than the original description of the high digestibility types, where an uncooked digestibility of 10–15% improvement was reported²³. With all the sorghums, malting improved protein digestibility on average 25%. This can be attributed primarily to

modification of the sorghum endosperm protein body and protein matrix by the endogenous proteases during germination²¹. Like the raw grain, the malts of the HPDS also had higher protein digestibility than the NPDS. However, the percentage increase in protein digestibility with malting in HPDS was considerably less than with the NPDS. This indicates that the protein modification which occurred during malting had a greater effect on protein digestibility than the high protein digestibility trait.

Table II. Effects of malting and mashing with normal protein digestibility sorghum cultivars (NPDS) and high protein digestibility sorghum (HPDS) cultivars on FAN (mg/100 g sorghum)^a.

Sorghum cultivar	Grain FAN	Malt FAN	FAN produced during malting ^b	FAN at end of mashing	FAN produced during mashing ²
HPDS parent	12.1 ² ± 0.4	83.3 ³ ± 2.4	71.2	96.1 ³ ± 3.3	12.8
HPDS progeny	15.5 ⁴ ± 0.7	103.4 ⁴ ± 2.6	87.9	120.1 ⁴ ± 2.4	16.7
Means HPDS	13.8	93.4	79.6 (577)	108.1	14.8 (16)
NPDS parent	10.6 ¹ ± 0.8	73.1 ² ± 2.5	62.5	83.8 ² ± 2.2	10.8
NPDS control	14.2 ^{2,3} ± 0.7	50.4 ¹ ± 1.5	36.2	57.6 ¹ ± 0.8	7.2
Means NPDS	12.4	61.8	49.4 (398)	70.7	9.0 (15)
Grand Means	13.1	77.6	64.5	89.4	11.9

^a All results are means of two experiments with standard deviations. Means with different number superscripts within the columns differ significantly (p<0.05).

^b Percentage increase in parentheses.

Table III. Effect of exogenous proteases on FAN production when mashing with unmalted normal protein digestibility sorghum cultivars (NPDS) and high protein digestibility sorghum (HPDS) cultivars (mg/100 g sorghum)^a.

Sorghum cultivar	FAN at end of mashing		FAN produced ^b	
	Flavourzyme	Bioprotease	Flavourzyme	Bioprotease
HPDS parent	51.2 ² ± 1.3	80.6 ³ ± 1.7	39.1	68.5
HPDS progeny	53.4 ² ± 2.4	89.9 ⁴ ± 1.1	37.9	74.4
Mean HPDS	52.3	83.8	38.5 (279)	71.5 (518)
NPDS parent	45.0 ¹ ± 2.2	63.5 ¹ ± 2.1	34.4	52.9
NPDS control	44.3 ¹ ± 1.0	67.8 ² ± 2.5	30.1	53.5
Mean NPDS	44.7	65.7	32.3 (260)	53.2 (429)
Grand means	48.5 ± 4.5	75.5 ± 12.1	35.4	62.3

¹ All results are means of two experiments with standard deviations. Means with different number superscripts within the columns differ significantly (p<0.05).

² Percentage increase in parentheses compared to the average FAN levels in the grains.

Table IV. Effect of exogenous proteases on FAN production when mashing with malted normal protein digestibility sorghum cultivars (NPDS) and high protein digestibility sorghum (HPDS) cultivars (mg/100 g malt)^a.

Sorghum cultivar	FAN at end of mashing		FAN produced ^b	
	Flavourzyme	Bioprotease	Flavourzyme	Bioprotease
HPDS parent	112.7 ³ ± 1.9	148.1 ² ± 4.1	29.4	64.9
HPDS progeny	129.0 ⁴ ± 1.2	174.4 ³ ± 2.6	25.6	71.0
Mean HPDS	120.9	161.3	27.5 (29)	68.0 (73)
NPDS parent	91.9 ² ± 1.8	135.1 ² ± 4.1	18.8	62.0
NPDS control	69.5 ¹ ± 2.4	97.6 ¹ ± 2.1	19.1	47.2
Mean NPDS	80.7	116.4	19.0 (31)	54.6 (88)
Grand Means	100.8 ± 25.8	138.8 ± 32.0	23.2	61.2

^a All results are means of two experiments with standard deviations in brackets. Means with different number superscripts within the columns differ significantly (p<0.05).

^b Percentage increase in parentheses compared to the average FAN levels in the malts.

The level of FAN in the HPDS raw grains was similar to that in the NPDS (Table II). This indicates that despite the higher lysine content and digestibility of these HPDS, the proportion of nitrogen as protein, as opposed to amino acids and peptides, was the same as in the NPDS. As expected, there was a very large increase in FAN, on average 65 mg/100 g sorghum in all the sorghums with malting, which was as a result of the action of the endogenous proteolytic enzymes^{6,11,14}. The malts of the HPDS contained substantially higher levels of FAN than those of the NPDS, on average 32 mg FAN /100 g (51%) more. This was presumably because of the higher digestibility of the protein bodies (Fig. 2). However, the quantity of FAN produced when all sorghum malt types were mashed was low, on average only 12 mg/100 g sorghum, in line with earlier reports^{7,19}. Further, with the HPDS malts, FAN

production during mashing was only slightly higher than that from the NPDS, on average 6 mg/100 g more. To investigate this further, both raw grain and malt of HPDS and NPDS were mashed with exogenous protease enzymes.

With both HPDS and NPDS raw grain there was greater FAN production during mashing with Bioprotease P Conc than with Flavourzyme, on average 62 and 35 mg/100 g sorghum (Table III). This can be attributed to the greater activity of the Bioprotease P Conc enzyme preparation (see under Proteases). However, the difference in FAN production between the HPDS and NPDS was low, on average 18 and 6 mg/100 g sorghum, for Bioprotease P Conc and Flavourzyme respectively. The same was found when malts of the HPDS and NPDS were mashed with the exogenous protease enzymes (Table IV).

In this case, the difference in FAN production was 13 and 9 mg/100 g, respectively. These results are in agreement with those for sorghum malts mashed on their own (Table II) and show that under a wide range of conditions FAN production from HPDS during mashing was only slightly higher than from NPDS. It is probable that the wet-heat treatment of mashing still caused intermolecular disulphide-bonded cross-linking of the kafirin proteins^{4,12} in these HPDS, which reduces the susceptibility of kafirin to hydrolysis³.

Notwithstanding this, Table IV shows that the combination of using HPDS, malting these sorghums and applying exogenous proteases resulted in substantial wort FAN being obtained from sorghum, in this case 121–161 mg/100 g sorghum. Such levels meet the recommended adequate wort FAN levels of 130–150 mg/L for support of optimal yeast growth and rapid fermentation^{13,17}.

CONCLUSIONS

HPDS have substantially higher protein digestibility compared to NPDS, when measured by the pepsin (endo-protease-type) assay. They produce malt with substantially higher levels of FAN. However, they yield only slightly higher FAN when mashed. This is probably as a result of wet-heat induced cross-linking of the kafirin proteins, which reduces their susceptibility to proteolysis.

Notwithstanding this, HPDS could be very useful for improving FAN levels in sorghum brewing if they are malted. Also, to further increase FAN production when mashing with HPDS malts, it is recommended that the various lines are screened to select those with the highest protease activity.

ACKNOWLEDGEMENTS

The International Sorghum, Millet and Other Grains Collaborative Research Support Program (INTSORMIL) for financial support; Sun Grant Program, US Department of Transportation, Project No. DTOS59-07-G-00056 to D.B. Hays for financial support; Novozymes SA, Mr J. van Aswegen, Mr I. Kennedy and Mr B. Higgins, for technical and financial support; and Mr B. Cotterell and Mrs L. da Silva for their technical contributions.

LITERATURE CITED

1. AACC International, Crude Protein – Combustion Method, Method 46-30. Approved Methods of the American Association of Cereal Chemists, 10th ed., The Association: St. Paul, MN, 2000.
2. Bidlingmeyer, B. A., Cohen, S. A. and Tarvin, T. L., Rapid analysis of amino acids using pre-column derivatization. *J. Chromatog.*, 1984, **336**, 93-104.
3. Duodu, K. G., Taylor, J. R. N., Belton, P. S. and Hamaker, B. R., Mini review: Factors affecting sorghum protein digestibility. *J. Cereal Sci.*, 2003, **38**, 117-131.
4. Emmambux, M. N. and Taylor, J. R. N., Properties of heat-treated sorghum and maize meals and their prolamin proteins. *J. Agric. Food Chem.*, 2009, **57**, 1045-1050.
5. European Brewery Convention, Free Amino Nitrogen of Malt by Spectrophotometry, Method 4.10. Analytica-EBC, 5th ed., Fachverlag Hans Carl: Nuremberg, 1998.
6. Evans, D. J. and Taylor, J. R. N., Influence of cultivar and germination conditions on proteolytic activities in sorghum malt. *J. Inst. Brew.*, 1990, **96**, 399-402.
7. Goode, D. L., Halbert, C. and Arendt, E. K., Optimisation of mashing conditions when mashing with unmalted sorghum and commercial enzymes. *J. Am. Soc. Brew. Chem.*, 2003, **61**, 69-78.
8. Hamaker, B. R., Kirleis, A. W., Butler, L. G., Axtell, J. D. and Mertz, E. T., Improving the in vitro protein digestibility of sorghum with reducing agents. *Proc. Nat. Acad. Sci. USA*, 1987, **84**, 626-628.
9. International Association for Cereal Science and Technology (ICC), ICC Draft Standard No. 174, Determination of Germinative Energy of Sorghum Grain, ICC: Vienna, 2009.
10. Mackintosh, I. and Higgins, B., The development of a sorghum based lager beer in Uganda: A model of co-operation between industry and government in the development of local ingredients for the production of quality lager beer and consequential benefits for the parties involved. *Aspects Appl. Biol.*, 2004, **72**, 235-245.
11. Mokhawa, G., Ralefala, M. P., Bulawayo, B. B. and Ezeogu, L. I., Qualitative zymographic studies of major proteinases from malted sorghum (*Sorghum bicolor* L.) and effects of cultivar. *Cereal Chem.*, 2009, **86**, 575-581.
12. Ng'andwe, C. C., Hall, A. N. and Taylor, J. R. N., Proteolysis of sorghum endosperm proteins when mashing with raw grain plus exogenous protease and potassium metabisulphite. *J. Inst. Brew.*, 2008, **114**, 343-348.
13. O'Connor-Cox, E. S. C. and Ingledew, W. M., Wort nitrogenous sources- Their use by brewing yeasts: A review. *J. Am. Soc. Brew. Chem.*, 1989, **47**, 102-108.
14. Ogbonna, A. C., Obi, S. K. C. and Okolo, B. N., Optimization of proteolytic activities in malting sorghum. *Process Biochem.*, 2004, **39**, 711-716.
15. Oria, M. P., Hamaker, B. R., Axtell, J. D. and Huang, C.-P., A highly digestible sorghum mutant cultivar exhibits a unique folded structure of endosperm protein bodies. *Proc. Nat. Acad. Sci. USA*, 2000, **97**, 5065-5070.
16. Oria, M. P., Hamaker, B. R., Axtell, J. D. and Shull, J. M., Resistance of sorghum α -, β -, and γ -kafirins to pepsin digestion. *J. Agric. Food Chem.* 1995, **43**, 2148-1160.
17. Pickerell, A. T. W., The influence of free alpha-amino nitrogen in sorghum beer fermentations. *J. Inst. Brew.*, 1986, **92**, 568-571.
18. Rooney L. W. and Miller F. R., Variation in the kernel structure and kernel characteristics of sorghum. Proceedings of the International Symposium on Sorghum Grain Quality. International Crops Research Institute for the Semi-Arid Tropics: Patancheru, India, 1982, pp. 143-162.
19. Taylor, J. R. N. and Boyd, H. K., Free α -amino nitrogen production in sorghum beer mashing. *J. Sci. Food Agric.*, 1986, **37**, 1109-1117.
20. Taylor, J. R. N., Dewar, J. and Joustra, S. M., Accepted Methods in Southern Africa of Sorghum Malting and Brewing Analyses, Method No. 6, Laboratory Malting of Sorghum. Sorghum Malting Technology Course and Methods of Sorghum Grain and Malt Quality Analysis, University of Pretoria: Pretoria, 2005.
21. Taylor, J. R. N., Novellie, L. and Liebenberg, N. v. d. W., Protein body degradation in the starchy endosperm of germinating sorghum. *J. Exp. Bot.*, 1985, **36**, 1287-1295.
22. Tesso, T., Ejeta, G., Chandrashekar, A., Huang, C.-P., Tandjung, A., Lewamy, M., Axtell, J. D., and Hamaker, B. R., A novel modified endosperm texture in a mutant high-protein digestibility/high-lysine sorghum (*Sorghum bicolor* (L.) Moench). *Cereal Chem.*, 2006, **83**, 194-201.
23. Weaver, C. A., Hamaker, B. R. and Axtell, J. D., Discovery of grain sorghum germ plasm with high uncooked and cooked in vitro protein digestibility. *Cereal Chem.*, 1998, **75**, 665-670.
24. Winn, J. A., Mason, E., Robbins, A. L., Rooney, W. L. and Hays, D. B., QTL mapping of a high protein digestibility trait in *Sorghum bicolor*. *Int. J. Plant Genomics* 2009. (doi:10.1155/2009/471853)

(Manuscript accepted for publication May 2011)