

Preventing Bloodstream Infections and Death in Zambian Neonates: Impact of a Low-cost Infection Control Bundle

Lawrence Mwananyanda,^{1,2} Cassandra Pierre,³ James Mwansa,^{4,5} Carter Cowden,⁶ A. Russell Localio,⁷ Monica L. Kapasa,⁸ Sylvia Machona,⁸ Chileshe Lukwesa Musyani,⁴ Moses M. Chilufya,¹ Gertrude Munanjala,¹ Angela Lyondo,⁴ Matthew A. Bates,⁹ Susan E. Coffin,^{6,a} and Davidson H. Hamer^{2,3,a}

¹Right to Care, Lusaka, Zambia; ²Department of Global Health, Boston University School of Public Health, and ³Section of Infectious Diseases, Department of Medicine, Boston Medical Center, Massachusetts; ⁴Department of Pathology and Microbiology, University Teaching Hospital, and ⁵Lusaka Apex Medical University, Zambia; ⁶Division of Infectious Diseases, Children's Hospital of Philadelphia, and ⁷Division of Biostatistics, Department of Biostatistics, Epidemiology and Informatics, University of Pennsylvania Perelman School of Medicine, Philadelphia; ⁸Neonatal Intensive Care Unit, University Teaching Hospital, Lusaka, Zambia; and ⁹School of Life Sciences, University of Lincoln, United Kingdom

(See the Editorial Commentary by Johnson and Milstone on pages 1368–9.)

Background. Sepsis is a leading cause of neonatal mortality in low-resource settings. As facility-based births become more common, the proportion of neonatal deaths due to hospital-onset sepsis has increased.

Methods. We conducted a prospective cohort study in a neonatal intensive care unit in Zambia where we implemented a multifaceted infection prevention and control (IPC) bundle consisting of IPC training, text message reminders, alcohol hand rub, enhanced environmental cleaning, and weekly bathing of babies ≥ 1.5 kg with 2% chlorhexidine gluconate. Hospital-associated sepsis, bloodstream infection (BSI), and mortality (>3 days after admission) outcome data were collected for 6 months prior to and 11 months after bundle implementation.

Results. Most enrolled neonates had a birth weight ≥ 1.5 kg (2131/2669 [79.8%]). Hospital-associated mortality was lower during the intervention than baseline period (18.0% vs 23.6%, respectively). Total mortality was lower in the intervention than prior periods. Half of enrolled neonates (50.4%) had suspected sepsis; 40.8% of cultures were positive. Most positive blood cultures yielded a pathogen (409/549 [74.5%]), predominantly *Klebsiella pneumoniae* (289/409 [70.1%]). The monthly rate and incidence density rate of suspected sepsis were lower in the intervention period for all birth weight categories, except babies weighing <1.0 kg. The rate of BSI with pathogen was also lower in the intervention than baseline period.

Conclusions. A simple IPC bundle can reduce sepsis and death in neonates hospitalized in high-risk, low-resource settings. Further research is needed to validate these findings in similar settings and to identify optimal implementation strategies for improvement and sustainability.

Clinical Trials Registration. NCT02386592.

Keywords. neonatal sepsis; *Klebsiella pneumoniae*; infection prevention; chlorhexidine; Zambia.

Approximately 2.7 million neonatal deaths occurred in 2015, accounting for 46% of all under-5 child mortality [1]. Almost all (99%) neonatal deaths occur in low- and middle-income countries (LMICs) and about one-fifth are attributable to neonatal sepsis and meningitis [2]. Bloodstream infections (BSIs), the most common hospital-associated infection in neonates, occur 3–20 times more frequently in LMICs than industrialized countries [3]. The excess burden of neonatal BSIs in LMICs is fueled by shortages of trained staff and basic resources, crowded clinical settings, malfunctioning equipment, increasing antimicrobial resistance, and suboptimal infection prevention and control (IPC) practices [3, 4].

Neonatal sepsis is a significant cause of morbidity and mortality in the largest referral neonatal intensive care unit (NICU) in Lusaka, Zambia (University Teaching Hospital [UTH]). A recent study in this NICU found 43% mortality among neonates with sepsis, predominantly due to multidrug-resistant *Klebsiella pneumoniae* [5]. IPC bundles composed of simple, relatively low-cost interventions [6, 7] might reduce the incidence of BSI and neonatal mortality. To assess the effectiveness of low-cost measures in LMIC settings, we determined the incidence of hospital-associated neonatal sepsis and studied the impact of a novel IPC bundle on hospital-associated BSI and mortality among neonates in the UTH NICU.

METHODS

We performed a prospective observational cohort study of neonates admitted to the NICU that included a 6-month baseline period (“baseline”), 6 weeks of bundle implementation (“implementation”), and 11 months of intervention assessment (“intervention”) (Supplementary Figure 1).

Received 15 September 2018; editorial decision 19 November 2018; accepted 24 December 2018; published online December 28, 2018.

^aS. E. C. and D. H. H. contributed equally to this work.

Correspondence: D. H. Hamer, Department of Global Health, Boston University School of Public Health, Crosstown 306, 801 Massachusetts Ave, Boston, MA 02118 (dhamer@bu.edu).

Clinical Infectious Diseases® 2019;69(8):1360–7

© The Author(s) 2018. Published by Oxford University Press for the Infectious Diseases Society of America. All rights reserved. For permissions, e-mail: journals.permissions@oup.com. DOI: 10.1093/cid/ciy1114

The study site, UTH, is a large tertiary care public hospital in Lusaka, Zambia, which is typically over census (60 cots and an average daily census of 75 infants). There are approximately 3300 admissions per year. The average patient-to-nurse ratio is 20:1 ([Supplementary Appendix A](#) contains additional information about the study site).

On weekdays from 8 AM to 5 PM, study staff enrolled neonates admitted to the NICU from September 2015 through March 2017. For neonates admitted during evenings and weekends, enrollment was attempted the next working day. Neonates were excluded if their mothers were <18 years old unless a legal guardian ≥18 years of age gave consent for both mother and baby to participate.

The Boston University Medical Campus, Excellence in Research Ethics and Science Converge (Zambia), and the Children's Hospital of Philadelphia institutional review boards; the Zambian Ministry of Health Research Secretariat; and the UTH Department of Pediatrics reviewed and approved the study. Written informed consent was obtained from the neonate's mother or legal guardian in English or the most common local languages, Nyanja and Bemba. A data and safety monitoring board, consisting of 3 independent experts in pediatrics, newborn health, and clinical trials in LMICs, met twice during the course of the study; no concerns were identified.

Study Outcomes and Definitions

The primary outcome was all-cause mortality among neonates hospitalized at the UTH NICU for ≥3 days ("hospital-associated mortality") to ensure that enrolled babies were exposed to the NICU-based interventions. Secondary outcomes included incidence of hospital-associated (onset ≥3 days after NICU admission) suspected sepsis and BSIs.

Blood cultures were obtained on all neonates with clinically suspected sepsis, defined as fever or hypothermia, tachycardia or bradycardia, hypoglycemia, respiratory difficulty, new-onset seizures, lethargy, poor feeding, abdominal distention, vomiting, diarrhea, or poor perfusion.

Positive blood cultures were classified as either BSI with pathogen or BSI with contaminant organism using commonly accepted criteria [8, 9]. [Supplementary Appendix A](#) contains additional description of classification of infants who experienced multiple episodes of suspected sepsis.

IPC Interventions

Our IPC bundle included (1) IPC training; (2) locally manufactured alcohol hand rub; (3) weekly bathing with 2% aqueous chlorhexidine gluconate (CHG; Clorox Company, Oakland, California); (4) daily IPC reminders via short messaging service (SMS); and (5) enhanced routine cleaning of the environment including potential reservoirs of infection (eg, sinks and suction machines), with a focus on daily cleaning of high-touch surfaces and moving from clean to dirty. We

did not introduce any changes in the cleaning and disinfection products used.

The IPC training was developed based on Jhpiego's curriculum [10]. Two identical trainings were held over 2 consecutive half-days. Study team members (C. P., S. E. C., C. L. M., J. M., and M. A. B.) taught all modules. Breakout sessions focused on role-specific practices (eg, environmental disinfection for cleaners, CHG bathing for nurses).

Alcohol hand rub was made according to a World Health Organization (WHO)-recommended formulation for local production [11]. The NICU pharmacist oversaw production and ensured continual availability. Hand rub dispensers were either wall-mounted or on mobile stands. Two dispensers were positioned in strategic locations close to patient care in all rooms. Study personnel performed daily checks of dispensers and replaced empty containers as necessary. Posters promoting and demonstrating proper technique for the WHO Five Moments of Hand Hygiene were prominently displayed.

All enrolled neonates admitted during the implementation and intervention periods underwent CHG bathing (sparing head and face) at admission and weekly thereafter unless they met an exclusion criteria: severe skin disease, an open wound, birth weight <1.5 kg, or estimated gestational age <32 weeks. Once newborns achieved both a weight of ≥1.5 kg and corrected gestational age of ≥32 weeks, they were eligible for CHG bathing, criteria currently used by many NICUs in the United States [12].

During the intervention period, daily SMS messages designed to reinforce IPC practices and concepts taught in the training sessions were sent to all NICU staff once daily. The messages included reminders on hand hygiene, appropriate glove use, and peripheral catheter care, and urged staff to reassess the duration and need for indwelling catheters. Additional details of the implementation of these interventions are provided in [Supplementary Appendix B](#).

Data Collection

Prospective Data

At enrollment, study staff interviewed mothers and reviewed maternal medical records to identify perinatal exposures, such as human immunodeficiency virus (HIV) status and prenatal infections. No observational data were collected on hand hygiene, environmental cleaning, or other IPC practices due to limited study staffing.

Historical Data

Using multiple sources, we assembled a limited data set to describe comprehensive neonatal mortality at the study NICU prior to study initiation. Limited data on all admitted neonates from October 2014 to March 2017 were extracted from the unit's discharge log, which was maintained by the unit clerk and completed at the time of each baby's discharge or death. The log

included dates of admission and discharge, birth weight, admission diagnoses, and vital status at discharge. Additionally, we captured similar data from a prior study's database from June 2013 to September 2014 [5]. See [Supplementary Appendix A](#) for additional description of analysis of historical data.

Microbiologic Data

One blood culture was obtained on all neonates with clinically suspected sepsis at the time of symptom onset. Urine and cerebrospinal fluid cultures were rarely obtained. Typically, a second blood culture was obtained if a patient developed a new episode of suspected sepsis. Pediatric blood culture bottles were inoculated with .5–1.0 mL of blood and incubated in the BD Bactec blood culture system (BD Life Sciences, Franklin Hills, New Jersey). Organisms isolated were tested for antibiotic susceptibilities using the VITEK Compact 2 identification and antibiotic susceptibility test automated machine (bioMérieux, Durham, North Carolina).

Data Management and Analysis

Prospective data were collected by full-time study staff into an electronic data collection platform implemented on electronic tablets (SurveyCTO, Cambridge, Massachusetts). Data were downloaded into SAS version 9.4 (SAS Institute, Cary, North Carolina) and Stata version 15.1 (StataCorp, College Station, Texas) software on a secure server.

Our prespecified analytic plans included aggregating data by month and constructing 3 discrete time periods (baseline, implementation, and intervention) for descriptive and statistical comparisons. Enrolled neonates were characterized by maternal demographic and clinical characteristics, neonatal clinical characteristics, and outcomes. This study was registered at ClinicalTrials.gov (identifier NCT02386592).

Prospective Data

We evaluated the effect of the study intervention on hospital-associated mortality and secondary outcomes of hospital-associated suspected sepsis, BSI with pathogen, or BSI with contaminant.

We used a simple piecewise regression with time as the covariate of interest, and with 2 knots to reflect possible discontinuity in the rates at the start of the implementation (month 7) and intervention (month 9) periods. We examined the change in levels and trajectories of hospital-associated mortality as well as our secondary outcomes over time.

Monthly rates of mortality and BSI with pathogen were examined among study participants. Smoothed rates were also calculated using a standard smoother for exploratory data analysis to reduce noise [13].

Microbiologic Data

All blood cultures that grew 1 or more organisms were included in this analysis. Blood cultures that grew ≥ 2 organisms were

classified as polymicrobial; if both a pathogen and a common skin commensal (contaminant) were isolated from the same culture, it was designated as BSI with pathogen.

Sensitivity Analysis

We conducted a sensitivity analysis to determine the impact of unmeasured confounders on observed mortality using “E-values.” The E-value determines how large an unmeasured confounder must be to negate the observed effect of the intervention ([Supplementary Appendix B](#)) [14].

RESULTS

Of 3686 newborns screened, 3348 (90.8%) were enrolled but only 2669 were fully evaluable, primarily due to death prior to day 3 of hospitalization in the study NICU ([Figure 1](#)).

The median maternal age and proportion of HIV-infected mothers were similar in all 3 time periods ([Table 1](#)). There were no differences in newborn characteristics in the 3 time periods with the exception of birth weight. There were no adverse events associated with CHG bathing.

Hospital-associated Mortality

Overall hospital-associated mortality for enrolled neonates was 20.1% and was lower during the intervention (18.0%) than the baseline (23.6%) period. Hospital-associated mortality was lowest for neonates >2.5 kg and highest in neonates <1.0 kg ([Table 2](#)). Hospital-associated mortality declined from baseline to the intervention period in all birth weight categories.

Total Mortality

We compared the historic monthly mortality data to the total mortality (early and hospital associated) during the intervention period ([Figure 2A](#)) and observed reduced rates of total mortality during the intervention period. When stratified by calendar month, the absolute mean monthly mortality reduction was -9.1 (95% confidence interval [CI], -10.8 to -7.3) percentage points, and the overall reduction in relative mortality was 21.0% (relative risk [RR], .79 [95% CI, .76–.83]). Heterogeneity of mortality reduction by calendar month was limited, with relative mortality ranging from .64 to .95 during the intervention period.

Suspected Sepsis and Bloodstream Infection

Half of the enrolled neonates (1344/2669 [50.4%]) experienced 1 or more episodes of suspected sepsis. The monthly rate of suspected sepsis was lower in the intervention than baseline period ([Figure 2B](#)). Similarly, incidence density rates for suspected sepsis were lower during the intervention than baseline period for all birth weight categories, except babies weighing <1.0 kg ([Table 3](#)).

Less than half of episodes of suspected sepsis (549/1344 [40.8%]) were associated with a positive blood culture, and the




Figure 1. Assembly of study cohort. ^aUnavailable to provide consent due to early death or discharge. ^bDied prior to neonatal intensive care unit day 3 (n = 670) or missing study maternal–neonatal and/or completion form (n = 9).

proportion of neonates with a positive blood culture was inversely associated with birth weight category. Pathogens were isolated from most positive blood cultures (409/549 [74.5%]). The most frequently isolated pathogens were *K. pneumoniae* (289/409 [70.1%]) and *Enterococcus* species (70/409 [18.6%]). The rate of BSI with pathogen was higher for babies of all birth weights in the baseline than the intervention period (Figure 3). Similarly, the incidence density rate of BSI with pathogen was higher for all birth weight categories during the baseline than the intervention period, except for neonates <1.0 kg (Table 3).

The daily hazard of death and BSI varied by duration of NICU stay and study period (Supplementary Figure 2). The daily hazard of death was lower during the first week of hospitalization in the intervention than baseline period. In contrast, the daily hazard of BSI with pathogen was highest on NICU day 3 (first day of study observation) for all babies, but lower for babies hospitalized during the intervention than the baseline period.

As a sensitivity analysis, we estimated the magnitude of unmeasured confounding, not already accounted for by our measured factors and design, needed to negate our observed association (RR, .79) between our intervention and hospital-associated

mortality. Such a confounder would need to be associated with our intervention and mortality by the same relative risk of ≥ 1.85 , and not lower. A confounder with less strong associations could not explain away this mortality reduction. We cannot identify such a strong unmeasured factor or a differential exposure that might account for the observed association. Additional sensitivity analyses, their interpretation, and the reference for the calculations are shown in Supplementary Appendix C.

DISCUSSION

There were significant reductions in the rates of suspected sepsis, BSI, and mortality among neonates in a low-resource NICU following implementation of a low-cost IPC bundle. We found lower rates of total and hospital-associated neonatal mortality after implementation of the IPC bundle. Additionally, we observed that hospital-associated mortality decreased across all birth weight groups, although the reductions were greatest among neonates <2.5 kg. These findings are similar to observations made in a study performed in 2 NICUs in Manila, Philippines, where a simpler bundle (composed of introduction of locally produced alcohol-based hand rub, IPC training, and checklists) reduced overall neonatal mortality [6].

Table 1. Characteristics of Enrolled Mothers and Neonates in the Neonatal Intensive Care Unit (N = 2669)

Characteristic	Baseline (n = 852)	Implementation (n = 268)	Intervention (n = 1549)	Significance ^a
Maternal characteristics				
Median age, y (IQR)	25 (21–31)	24 (20–31)	25 (21–31)	<i>P</i> = .44
HIV status				<i>P</i> = .83
Infected	167 (20)	59 (22)	320 (21)	
Uninfected	622 (74)	207 (78)	1219 (79)	
Unknown	63 (7)	2 (0.7)	6 (0.4)	
Neonatal characteristics				
Birth weight				<i>P</i> < .001
<1.0 kg	12 (1)	6 (2)	48 (3)	
1–1.49 kg	141 (17)	40 (15)	291 (19)	
1.5–2.49 kg	249 (29)	92 (34)	516 (33)	
≥2.5 kg	450 (53)	130 (49)	694 (45)	
Mode of delivery				<i>P</i> = .92
Spontaneous vaginal delivery	648 (77)	198 (74)	1162 (75)	
Spontaneous vaginal delivery with instrumentation	9 (1)	8 (3)	51 (3)	
Cesarean delivery	189 (22)	61 (23)	330 (21)	
Unknown	6 (0.7)	1 (0.4)	6 (0.4)	

Data are presented as no. (%) unless otherwise indicated.

Abbreviations: HIV, human immunodeficiency virus; IQR, interquartile range.

^aComparing baseline to intervention period.

We observed reductions in the rates of sepsis and BSI with pathogen across all birth weight categories except neonates weighing <1.0 kg. Early deaths due to extreme prematurity and difficulties obtaining a sample for blood culture might have contributed to this finding. Prior quasi-experimental studies of the capacity of a simple IPC bundle to prevent neonatal sepsis have yielded inconsistent results [6, 15]. Our data suggest that the observed mortality reductions were potentially achieved by reducing mortality among all neonates (both those who did and did not experience sepsis), as well as by reducing sepsis. Our crude mortality analysis suggests that our intervention may have dampened the apparent seasonal variation with elimination of the peaks associated with the dry season. Although the mechanism by which this change might have been exerted is unclear, this finding deserves future study.

Our low-cost IPC bundle included both evidence-based and novel interventions. Practices such as staff education, hand hygiene, and environmental cleaning are well-recognized facets of

IPC practice in both resource-advantaged and resource-limited settings [16, 17]. In contrast, few published studies from LMIC hospitals have described the use of CHG bathing as a strategy to prevent sepsis or death. Studies performed in South Africa and Pakistan found that CHG bathing was not associated with a reduced rate of neonatal sepsis or death, although the concentration of CHG used in these studies was lower than used in our study (2% vs <1%) [18, 19]. Studies from North America have described the impact of daily CHG bathing on rates of bacteremia and infection with multidrug-resistant organisms in children and adult patients with central venous catheters [20, 21]. Among nonhospitalized neonates, the use of CHG for cord care is associated with reduced rates of omphalitis, neonatal sepsis, and death in some, but not all, settings [22–26]. Although reported in other studies, we did not observe adverse events (such as skin irritation) in neonates exposed to CHG [27].

While numerous studies have described the impact of a single intervention (eg, staff education or introduction of

Table 2. Hospital-associated Mortality, by Birth Weight Category

Birth Weight Category	Baseline			Implementation			Intervention			Total		
	No.	No. Died	%	No.	No. Died	%	No.	No. Died	%	No.	No. Died	%
<1 kg	10	9	90.0	6	5	83.3	46	36	78.3	62	45	80.4
1–1.49 kg	141	74	52.5	40	22	55.0	291	89	30.6	472	185	39.2
1.5–2.49 kg	249	55	22.1	92	10	10.9	516	69	13.4	857	134	15.6
≥2.5 kg	450	62	13.8	130	19	14.6	694	85	12.3	1274	166	13.0
Total	852	201	23.6	268	56	20.1	1549	279	18.0	2669	536	20.1




Figure 2. A, Overall (early and hospital-associated) crude mortality rate of patient-days at risk, grouped by month of admission to the neonatal intensive care unit (NICU) (dots). Smoothed mortality rates (line) reflect standard smoothing for exploratory data analysis. Seasonal variation was pronounced prior to the intervention compared with after the intervention. B, Crude monthly rate of suspected sepsis (dots) and smoothed monthly rate (line) of patient-days at risk, grouped by month of admission to the NICU among enrolled neonates admitted to the University Teaching Hospital NICU from September 2015 through March 2017.

alcohol hand rub) on colonization and neonatal sepsis, few studies have evaluated the impact of a bundle of IPC interventions. Prior reports have mainly focused on the control of outbreaks in LMIC NICUs and have often lacked detail on the interventions [28]. Additionally, few studies performed in NICUs have assessed the impact of IPC interventions on hospital-associated mortality.

Given evidence of impact of the interventions, the relative simplicity and potential for sustainability of our IPC bundle merit highlighting. Alcohol hand rub was produced locally through using relatively inexpensive reagents. CHG was externally procured (at low cost) for neonatal bathing. Daily IPC reminders via SMS using customized messages for different cadres of NICU clinical and nonclinical personnel may have enhanced adherence to specific elements of our IPC bundle, such as hand hygiene, at low cost. Similarly, mobile technology-based health interventions have shown improved health worker

behaviors for initiatives as diverse as acute respiratory infection and malaria case management [29, 30].

Our study design did not allow a rigorous analysis of the relative contribution of individual bundle components. Such an analysis would require larger numbers of patients and an extended study period. Future studies are needed to dissect whether more financially and labor-intensive interventions (such as CHG bathing) are necessary to achieve maximal benefit.

Strengths of our study include the availability of robust historical data, use of low-cost interventions with locally available supplies, ability to control for or at least identify alternative explanations for our findings, and longitudinal data collection over sufficient time to account for seasonal variation. There are also several important limitations. First, as a prospective observational, single-site study, our findings may not be easily generalizable. However, a randomized trial of unit-level interventions would require many NICUs to achieve an adequate sample size.

Table 3. Incidence Density Rate (per 100 Patient-days) of Suspected Sepsis and Bloodstream Infection Pathogen, by Birth Weight Category

Birth Weight Category	Incidence Density Rate (per 100 Patient-Days)		Incidence Rate Ratio (95% CI)	Significance
	Baseline	Intervention		
Suspected sepsis				
<1 kg	3.57	4.92	1.38 (.53–4.57)	<i>P</i> = .53
1–1.49 kg	7.09	3.42	.48 (.37–.62)	<i>P</i> < .001
1.5–2.49 kg	8.07	5.06	.63 (.51–.77)	<i>P</i> < .001
≥2.5 kg	9.15	5.95	.65 (.55–.76)	<i>P</i> < .001
BSI due to pathogen				
<1 kg	2.86	1.76	.62 (.18–2.69)	<i>P</i> = .42
1–1.49 kg	3.74	1.03	.28 (.18–.42)	<i>P</i> < .001
1.5–2.49 kg	3.27	1.25	.38 (.26–.55)	<i>P</i> < .001
≥2.5 kg	3.02	1.43	.47 (.35–.65)	<i>P</i> < .001

Abbreviations: BSI, bloodstream infection; CI, confidence interval.




Figure 3. Proportion of enrolled neonates without sepsis, with culture-negative sepsis, bloodstream infection due to pathogen (BSI-pathogen), and bloodstream infection due to possible contaminant (BSI-contam), stratified by intervention and birth weight.

This was beyond the scope of our funding. Second, we were unable to enroll participants on weekends and during public holidays due to limited staff availability, although we have no evidence that weekend/holiday admissions fare differently than those enrolled in our study. Similarly, with available research personnel we were unable to capture detailed data on babies who died or were discharged within 3 days of NICU admission. Third, adherence to study interventions, including hand hygiene and environmental cleaning, could not be systematically assessed. This study was pragmatic, and as such generalizes to typical settings that might face imperfect adherence. Finally, we were unable to assess the sustainability of the observed benefits of the interventions. Although our interventions were selected based upon the relative low cost and ease of implementation, they did require supplemental resources such as dedicated effort from the hospital pharmacy (to produce and distribute the alcohol-based hand rub and CHG) and modest costs associated with purchase of CHG and use of SMS reminders. As future studies seek to reproduce these findings, they should assess the costs associated with these interventions. It will also be essential to determine how these individual interventions can be embedded in hospital operations and funded through routine budgeting to ensure sustainability.

In conclusion, we observed a reduction in neonatal sepsis and death associated with the implementation of an IPC bundle. Many

NICUs and special care nurseries in resource-limited settings have serious limitations in space, staffing, and funding that challenge implementation of many traditional IPC measures, such as cohorting, isolation, and surveillance cultures. Thus, low-cost bundled interventions may represent the most effective and sustainable measures to prevent neonatal sepsis and death in these settings.

Supplementary Data

Supplementary materials are available at *Clinical Infectious Diseases* online. Consisting of data provided by the authors to benefit the reader, the posted materials are not copyedited and are the sole responsibility of the authors, so questions or comments should be addressed to the corresponding author.

Notes

Acknowledgments. The authors thank the following for their kind support of the study: Surabhi Bhat, Kenya Murray, Nellisiwe Chizuni; the staff of the University Teaching Hospital neonatal intensive care unit (UTH NICU); Mr Musamali, the UTH NICU pharmacist; members of the data and safety monitoring board (Drs Theoklis Zaoutis, Nanthilile Mugala, and Albert Manasyan); and Dr Sylvester Sinyangwe, the Chair of the UTH Department of Paediatrics.

Disclaimer. The study was designed, analyzed, and implemented by the authors. The funder had no role in data collection, data analysis, or interpretation of the findings.

Financial support. This work was supported by the Thrasher Research Fund (grant number 12036).

Potential conflicts of interest. S. E. C. serves on a data safety and monitoring board for Merck & Co. All other authors report no potential conflicts. All authors have submitted the ICMJE Form for Disclosure of Potential

Conflicts of Interest. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed.

References

1. Liu L, Oza S, Hogan D, et al. Global, regional, and national causes of under-5 mortality in 2000-15: an updated systematic analysis with implications for the sustainable development goals. *Lancet* **2016**; 388:3027-35.
2. Lawn JE, Cousens S, Zupan J; Lancet Neonatal Survival Steering Team. 4 million neonatal deaths: When? Where? Why? *Lancet* **2005**; 365:891-900.
3. Zaidi AK, Huskins WC, Thaver D, Bhutta ZA, Abbas Z, Goldmann DA. Hospital-acquired neonatal infections in developing countries. *Lancet* **2005**; 365:1175-88.
4. Allegranzi B, Bagheri Nejad S, Boubesque C, et al. Burden of endemic health-care-associated infection in developing countries: systematic review and meta-analysis. *Lancet* **2011**; 377:228-41.
5. Kabwe M, Tembo J, Chilukutu L, et al. Etiology, antibiotic resistance and risk factors for neonatal sepsis in a large referral center in Zambia. *Pediatr Infect Dis J* **2016**; 35:e191-8.
6. Gill CJ, Mantaring JB, Macleod WB, et al. Impact of enhanced infection control at 2 neonatal intensive care units in the Philippines. *Clin Infect Dis* **2009**; 48:13-21.
7. Landre-Peigne C, Ka AS, Peigne V, Bougère C, Seye MN, Imbert P. Efficacy of an infection control programme in reducing nosocomial bloodstream infections in a Senegalese neonatal unit. *J Hosp Infect* **2011**; 79:161-5.
8. Hossain B, Islam MS, Rahman A, et al. Understanding bacterial isolates in blood culture and approaches used to define bacteria as contaminants: a literature review. *Pediatr Infect Dis J* **2016**; 35:S45-51.
9. Hossain B, Weber MW, Hamer DH, et al. Classification of blood culture isolates into contaminants and pathogens on the basis of clinical and laboratory data. *Pediatr Infect Dis J* **2016**; 35:S52-4.
10. Tietjen L, Bossemeyer D, McIntosh N. Infection prevention: guidelines for health-care facilities with limited resources. **2003**. Available at: <https://www.popline.org/node/276510>. Accessed 14 July 2018.
11. World Health Organization. WHO guidelines on hand hygiene in health care. Geneva, Switzerland: WHO, **2009**.
12. Johnson J, Bracken R, Tamma PD, Aucott SW, Bearer C, Milstone AM. Trends in chlorhexidine use in US neonatal intensive care units: results from a follow-up national survey. *Infect Control Hosp Epidemiol* **2016**; 37:1116-8.
13. Hoaglin DC, John W. Tukey and data analysis. *Stat Sci* **2003**; 18:311-8.
14. VanderWeele TJ, Ding P. Sensitivity analysis in observational research: introducing the E-value. *Ann Intern Med* **2017**; 167:268-74.
15. Murni I, Duke T, Kinney S, Daley A, Soenarto Y. Reducing hospital-acquired infections and improving the rational use of antibiotics in a developing country: an effectiveness study. *Arch Dis Childhood* **2015**; 100:454-9.
16. Pittet D, Allegranzi B, Boyce J; World Health Organization World Alliance for Patient Safety First Global Patient Safety Challenge Core Group of Experts. The World Health Organization guidelines on hand hygiene in health care and their consensus recommendations. *Infect Control Hosp Epidemiol* **2009**; 30:611-22.
17. Yokoe DS, Anderson DJ, Berenholtz SM, et al; Society for Healthcare Epidemiology of America. A compendium of strategies to prevent healthcare-associated infections in acute care hospitals: 2014 updates. *Infect Control Hosp Epidemiol* **2014**; 35:967-77.
18. Cutland CL, Madhi SA, Zell ER, et al; PoPS Trial Team. Chlorhexidine maternal-vaginal and neonate body wipes in sepsis and vertical transmission of pathogenic bacteria in South Africa: a randomised, controlled trial. *Lancet* **2009**; 374:1909-16.
19. Saleem S, Rouse DJ, McClure EM, et al. Chlorhexidine vaginal and infant wipes to reduce perinatal mortality and morbidity: a randomized controlled trial. *Obstet Gynecol* **2010**; 115:1225-32.
20. Milstone AM, Elward A, Song X, et al; Pediatric SCRUB Trial Study Group. Daily chlorhexidine bathing to reduce bacteraemia in critically ill children: a multi-centre, cluster-randomised, crossover trial. *Lancet* **2013**; 381:1099-106.
21. Huang SS, Septimus E, Kleinman K, et al; CDC Prevention Epicenters Program; AHRQ DECIDE Network and Healthcare-Associated Infections Program. Targeted versus universal decolonization to prevent ICU infection. *N Engl J Med* **2013**; 368:2255-65.
22. Mullany LC, Darmstadt GL, Khatry SK, et al. Topical applications of chlorhexidine to the umbilical cord for prevention of omphalitis and neonatal mortality in southern Nepal: a community-based, cluster-randomised trial. *Lancet* **2006**; 367:910-8.
23. Arifeen SE, Mullany LC, Shah R, et al. The effect of cord cleansing with chlorhexidine on neonatal mortality in rural Bangladesh: a community-based, cluster-randomised trial. *Lancet* **2012**; 379:1022-8.
24. Soofi S, Cousens S, Imdad A, Bhutto N, Ali N, Bhutta ZA. Topical application of chlorhexidine to neonatal umbilical cords for prevention of omphalitis and neonatal mortality in a rural district of Pakistan: a community-based, cluster-randomised trial. *Lancet* **2012**; 379:1029-36.
25. Sazawal S, Dhingra U, Ali SM, et al. Efficacy of chlorhexidine application to umbilical cord on neonatal mortality in Pemba, Tanzania: a community-based randomised controlled trial. *Lancet Glob Health* **2016**; 4:e837-44.
26. Semrau KEA, Herlihy J, Grogan C, et al. Effectiveness of 4% chlorhexidine umbilical cord care on neonatal mortality in Southern Province, Zambia (ZamCAT): a cluster-randomised controlled trial. *Lancet Global Health* **2016**; 4:e827-36.
27. Chapman AK, Aucott SW, Milstone AM. Safety of chlorhexidine gluconate used for skin antisepsis in the preterm infant. *J Perinatol* **2012**; 32:4-9.
28. Dramowski A, Aucamp M, Bekker A, Mehtar S. Infectious disease exposures and outbreaks at a South African neonatal unit with review of neonatal outbreak epidemiology in Africa. *Int J Infect Dis* **2017**; 57:79-85.
29. Chen Y, Yang K, Jing T, et al. Use of text messages to communicate clinical recommendations to health workers in rural China: a cluster-randomized trial. *Bull World Health Organ* **2014**; 92:474-81.
30. Zurovac D, Sudoi RK, Akhwale WS, et al. The effect of mobile phone text-message reminders on Kenyan health workers' adherence to malaria treatment guidelines: a cluster randomised trial. *Lancet* **2011**; 378:795-803.