

The antibacterial properties found in the Brassica Oleracea leaves. A Review

Top Journal Best Impact Factor

Related papers

[Download a PDF Pack](#) of the best related papers 



[Health-Affecting Compounds in Brassicaceae](#)

Robert Verporte

[The antimicrobial effects of glucosinolates and their respective enzymatic hydrolysis products on ba...](#)

Alfredo Carvalho Aires, Maria Saavedra

[Metal ion-inducing metabolite accumulation in Brassica rapa](#)

Ibrahim Abdel-farid

The antibacterial properties found in the Brassica Oleracea leaves. A Review.

DR D Ndayazi*, Pr ShimbilaKabangu**, Pr Yassa Y.Pierre***,
Pr Satheesh Kumar****

* Department of Surgery, Faculty of Medicine, University of Goma, Democratic Republic of Congo

** Department of Surgery, Faculty of Medicine and Pharmacy, University of Kisangani, University of Goma, Democratic Republic of Congo.

*** Catholic university Sapiensia of Goma, Democratic Republic of Congo

**** Lusaka Apex Medical University, Republic of Zambia

Corresponding Author: DR D Ndayazi

ABSTRACT : The antimicrobial activities have been searched previously on plant infection control and for food preservation. Current research have studied the mode of action of ITCs attracted attention of several researchers. The objective of this review is to discuss the current knowledge and hypothesis for the antibacterial effect of the leaves of the Brassica oleracea plant. (ITCs) of organic extract of leaves of the Brassica oleracea plant on some vegetative microorganism like *Candida albicans*, *Staphylococcus* spp., *Salmonella* spp., *Klebsiella* spp., *Bacillus* spp. and yeast.

Keys words: Brassica oleracea plant, antibacterial properties

Date of Submission: 04-04-2018

Date of acceptance: 19-04-2018

I. INTRODUCTION

Brassica oleracea is one the most vegetables consumed in the world and is found in local markets (Kusznierewicz et al., 2008). It is allied to the family of Brassicaceae (or Cruciferae) along with Collards, Brussels sprouts, Broccoli, Cauliflower and Kale (Jaiswal et al., 2011).

The above vegetable content an abundant phytochemicals such as flavonoids and glucosinolates and their hydrolysis products. It has a good proprieties that promote health and plays a big role in preventing against different conditions such us cancer, cancer, atherosclerosis, nephritis and diabetes mellitus (Taveira et al., 2009).

Moon (2011) in their study from a crude methanol extract prepared from fresh broccoli sprouts was extracted with hexane, chloroform, ethyl acetate, and butanol sequentially. Those substances had higher antibacterial actives us demonstrated in the same study in different dilutions used against the *Helicobacter pylori*.

The chloroform extract has shown greatest inhibition zones (>5 cm) against *Helicobacter pylori* s, followed by the hexane extract (5.03 cm), the ethyl acetate extract (4.90 cm), the botanical extract (3.10 cm), and the crude methanol extract (2.80 cm), however the residual water fraction did not demonstrate any inhibition zone (Moon et al., 2011). The priorities antibacterial were found among 18 sulforaphane and related compounds synthesized (6 amines, 6 isothiocyanates, and 6 nitriles), 2 amines, 6 isothiocyanates, and 1 Nitrile exhibited >5 cm inhibitory zones for *H. pylori* strain.

Researchers had conducted several studies to understand the antibacterial activity of cabbage juice that were observed with the glucosinolates degradation by-products found in the juice (Saeed, S. and P. Tariq, 2006; Stoewsand, G.S, 1995). The antibiotic and antifungal activity against a large number of bacteria, and used in the treatment of lung diseases (Chopra, D. and D. Simon, D, 2000, Dutta et al., 1998).

These researches have stimulated research to revisit the potential of plant products of different ingredients in the processing of other foods (Sebranek, J.G. and J. Bacus, 2007 and Honikel, 2008).

The purpose of this review article is to discuss the current knowledge and hypothesis for the antibacterial effect of the leaves of the Brassica oleracea plant.

II. METHODS

We conducted review articles using PubMed, embase, Google scholar and cochrane. The MesH word used were Brassica oleracea plant, antibacterial properties .

III. RESULTS AND DISCUSSION

ANTIMICROBIAL ACTIVITY OF THE PLANT EXTRACTS OF *BRASSICA OLERACEA* AGAINST SELECTED MICROBES.

Several studies were conducted by Swati et al., (2012) to discover the antimicrobial activity of the plant extract. Different concentration was used to the antimicrobial activity against three pathogenic microorganisms such as *Aspergillus fumigatus*, *Citrobacter diversions* and *Klebsiella pneumonia*. The maximum zone of inhibition was obtained for *Aspergillus fumigatus* and *Klebsiella pneumonia* at a concentration of 200µg/200µl. While *Klebsiella pneumonia* exhibited good sensitivity against both the concentrations and *Citrobacter* divergence showed medium sensitivity.

From the above studies, it is concluded that the traditional plants may represent new sources of anti-microbial with stable, biologically active components that can establish a scientific base for the use of plants in modern medicine.

Phytochemical analysis has studied the antimicrobial activities of the Brassica oleracea worldwide and the following substances were responsible for these activities against infections such as flavonoids, glycosides, saponins, steroids, terpenoids and alkaloids (Sibi et al., 2013).

The evidences in the literature have shown the bioactivity data obtained from the current study using the tested extracts of the plant demonstrated the potential to inhibit bacteria and fungi in the laboratory (Suganya. D et al., 2016).

The Laboratory sensitivity have shown *Pseudomonas aeruginosa* exhibited more inhibitory activity which represents the role of phytoconstituents towards the action of permeability on peptidoglycon layer.

The acetone extract component have demonstrate ability to create an acidity environment that lead to the disruption of the bacterial and fungal cell membrane. That made it to have a maximum antibacterial activity (Suganya. D et al., 2016)..

Phytochemical agents act as antimicrobial agent by inhibiting the extracellular enzyme acting on the substrates required for microbial growth or by inhibiting oxidative phosphorylation of microbial metabolism.

Several studies conducted found that active phytochemicals of *Brassica oleracea* had an antimicrobial activity against *Aspergillus fumigatus*, *Citrobacter diversens* and *Klebsiella pneumonia* (Swati Paul, 2012).

IV. CONCLUSION

Conclusion Traditional plants such as active phytochemicals of Brassica oleracea is found to have new sources of anti-microbial with the same proprieties that can be used in different medical interventions. However, further studies need to be conducted to assess the proprieties of the botanical preparation of those medicinal plants that can be extended in the field of pharmacology.

REFERENCE

- [1]. Jaiswal, A.K., Gupta, S., Abu-Ghannam, N. & Cox, S. (2011). Application of Baranyi function to model the antimicrobial properties of solvent extract from Irish York cabbage against food spoilage and pathogenic bacteria. *Food Science and Technology International*, 17(5), 495-502).
- [2]. Kusznierevicz B, Bartoszek A, Wolska L, Drzewiecki J, Gorinstein S and Namiesnik J. (2008). Partial characterization of white cabbages (*Brassica oleracea* var. capitata f. alba) from different regions by glucosinolates, bioactive compounds, total antioxidant activities and proteins. *LWT - Food Science and Technology* 41(1): 1–9.
- [3]. Taveira M, Pereira DM, Sousa C, Ferreres F, Andrade PB, Martins A, et al. (2009). In vitro cultures of *Brassica oleracea* l. var. costata dc: potential plant bioreactor for antioxidant phenolic compounds. *Journal of Agricultural and Food Chemistry* 57(4): 1247–1252.
- [4]. Saeed, S. and P. Tariq, 2006. Effects of some seasonal vegetables and fruits on the growth of bacteria. *Pakistan Journal of Biological Sciences*, 9(8): 1547-1551.
- [5]. Stoewsand, G.S, 1995. Bioactive organosulfur phytochemicals in *Brassica oleracea* vegetables. A review. *Food Chemistry and Toxicology*, (33):537-543.
- [6]. Chopra, D. and D. Simon, D, 2000. *The Chopra Centre Herbal Handbook*. Rider, London. Dutta, K., I. Rahman and K. Das, 1998. Antifungal activity of Indian plant extracts. *Mycoses*, (41):535-536.
- [7]. Sebranek, J.G. and J. Bacus, 2007. Cured meat products without direct addition of nitrate or nitrite: what are the issues? *Meat Science*, (77): 136–147.

- [8]. Honikel, K.-O, 2008. The use and control of nitrate and nitrite for the processing of meat products. *Meat Science*, (78): 68-76.
- [9]. Survey NS¹, Kumar B, Jang M, Yoon DY, Jung YS, Yang DC, Park SW. Two novel bioactive glucosinolates from Broccoli (*Brassica oleracea* L. var. *italica*) florets. *Bioorg Med Chem Lett*. 2012 Sep 1;22(17):5555-8. doi: 10.1016/j.bmcl.2012.07.016. Epub 2012 Jul 14.
- [10]. Moon JK¹, Kim JR, Ahn YJ, Shibamoto T. Analysis and anti-*Helicobacter* activity of sulforaphane and related compounds present in broccoli (*Brassica oleracea* L.) sprouts. *J Agric Food Chem*. 2010 Jun 9;58(11):6672-7.
- [11]. Ye XJ¹, Ng TB, Wu ZJ, Xie LH, Fang EF, Wong JH, Pan WL, Wing SS, Zhang YB.
- [12]. Protein from red cabbage (*Brassica oleracea*) seeds with antifungal, antibacterial, and anticancer activities. *J Agric Food Chem*. 2011 Sep 28;59(18):10232-8
- [13]. Swati Paul* Talha Bin Emran, Didyayyoti Saha, S.M. Zahid Hos. phytochemical and antimicrobial activity of the plant extracts of *brassicaoleracea* against selected microbes.. *Asian Journal of Pharmacy and Medical Science*. Vol 2 (2), 2012
- [14]. Suganya.D, Hussain Ali Fathima.M and Kanimozh.Ki. NTIBACTERIAL AND PHTYOCHEMICAL ANALYSIS ON *Brassica oleracea* var. *botrytis* Linn. *International Journal of Applied and Pure Science and Agriculture (IJAPSA)* Volume 02, Issue 03, [March - 2016

DR D Ndayazi "The antibacterial properties found in the Brassica Oleracea leaves. A Review." *IOSR Journal of Pharmacy (IOSRPHR)*, vol. 8, no. 4, 2018, pp. 82-84.